

Alyceclover Seed

Alyceclover Seed or Alyce Colver Seed (*Alysicarpus vaginalis*) is an annual, warm-season legume. It grows erect with spreading branches and may reach a height of 60" if not harvested. Although an annual, it can reestablish each year by natural reseeding if the crop is allowed to make seed in the fall. This may be difficult to accomplish since the crop is usually cut for hay or grazed closely, which eliminates seed production. Only seed of common alyceclover are available for planting. Common alyceclover is not a new crop, but has been grown in Florida for many years for either pasture or hay.

BUY "ALYCECLOVER" FOOD PLOT SEED

For orders over 250 lbs. please use our eQuote system for DISCOUNTS!

Site Selection

Alyceclover is adapted to well-drained soils with good fertility. These can be either upland sands or flatwoods with good drainage. Alyceclover does not perform well on soils that flood. Avoid sites with high levels of rootknot nematodes. These pests will attack common alyceclover and may severely damage the crop. Sites where alyceclover or other crops susceptible to rootknot nematodes have been grown in the previous season may have high levels of nematodes.

Seedbed Preparation

Alyceclover may be overseeded into a grass sod. The sod should be grazed or mowed closely to remove excess forage and cut with a medium-to-lightweight disk or chopper before seeding. The purpose of this cultivation is not to destroy the sod but to expose some soil so that seed can come in direct contact with it when planted. If a pasture drill is used, disking may be eliminated, but the sod must be closely grazed before planting.

Planting

Alyceclover should be planted between April 15 and June 30. If winter rains have been below normal and soil moisture is low in April and May, then overseeding of a grass sod should be delayed until the summer rains begin in June. Use 15 to 20 lb/A of seed. Use the high rate when the seedbed and soil condition are less favorable. The seed may be inoculated just prior to planting with N-fixing bacteria. Use the cowpea-type inoculant.

When planting into a grass sod that has been disked or chopped, the seed can be broadcast, followed by a harrow or drag of some type that will mix the seed with the soil, and then a cultipacker or roller may be useful to firm the soil. A good rain after broadcasting the seed will move the seed down and firm the soil, thus eliminating the need for harrowing and packing, but rainfall is unpredictable. When using a pasture drill, make sure the grass is cut or grazed short. When planting on a prepared seedbed, the seed are usually broadcast and covered with a cultipacker or planted with a grain drill. With either method, seed should be covered to a depth of 1/4 to 1/2". A well-prepared, level seedbed allows for planting at the right depth.

Alysicarpus vaginalis wildlife seed

Management and Utilization

Alyceclover planted early (April or May) will be tall enough for hay in mid to late August, but frequent rainfall may prevent harvesting at this time. Harvest can be delayed from 4 to 6 weeks without a severe loss in feeding value.

Alyceclover can be grazed after the plants are about 12" tall. The animals should be removed after the plants are grazed down in order to permit regrowth. Overgrazing could allow weeds to invade the stand. The first growth may be grazed down and animals removed and regrowth used for hay or seed production. Alyceclover can be used in a creep graze system for calves. Four-month-old calves have gained an additional 0.3 to 0.5 lb/day until they were weaned when alyceclover was available in a creep pasture.

Seed yields of up to 600 lb/A may be obtained. Seed may be combined directly or the plants may be mowed, cured in the swath, and then threshed with a combine with a pick-up attachment. The latter method will usually result in a larger amount of seed being saved. Harvest of seed should begin when about 1/2 of the seed pods are brown. This will usually occur about mid-October.

Courtesy of <http://edis.ifas.ufl.edu/>.